Noname manuscript No.
(will be inserted by the editor)

Lame equation in the algebraic form

Yoon Seok Choun

arXiv:1303.0873v5 [math-ph] 22 May 2013

Received: date / Accepted: date

Abstract Lame equation arises from deriving Laplace equation in ellipsoidal coordinates; in other words, it's called ellipsoidal harmonic equation. Lame functions are applicable to diverse areas such as boundary value problems in ellipsoidal geometry, chaotic Hamiltonian systems, the theory of Bose-Einstein condensates, etc.

In this paper I will apply three term recurrence formula[1] to the power series expansion in closed forms of Lame function in the algebraic form(infinite series and polynomial) and its integral forms including all higher terms of A_n's. I will show how to transform power series expansion of Lame function to an integral formalism mathematically for cases of infinite series and polynomial. One interesting observation resulting from the calculations is the fact that a ${}_2F_1$ function recurs in each of sub-integral forms: the first sub-integral form contains zero term of A'_ns, the second one contains one term of A_n's, the third one contains two terms of A_n's, etc. Section 6 contains additional examples of application in Lame function.

This paper is 6th out of 10 in series "Special functions and three term recurrence formula (3TRF)". See section 7 for all the papers in the series. Previous paper in series deals with the power series expansion of Mathieu function and its integral formalism [14]. The next paper in the series describes the power series and integral forms of Lame equation in the Weierstrass's form and its asymptotic behaviors[16].

Keywords Lame equation · Integral form · Three-term recurrence formula · Ellipsoidal harmonic function

Mathematics Subject Classification (2010) 33E05 · 33E10 · 34A25 · 34A30

Baruch College, The City University of New York, Natural Science Department, A506, 17 Lexington Avenue, New York, NY 10010
Tel.: +646-660-6200
Fax: + 646-660-6201
E-mail: ychoun@gc.cuny.edu; yoon.choun@baruch.cuny.edu; ychoun@gmail.com

1 Introduction

The sphere is a geometrical perfect shape, the set of points which are all equidistant from its center (a fixed point) in three-dimensional space. In contrast, an ellipsoid is a imperfect one, a surface whose plane sections are all ellipses or circles; the set of points are not same distance from the center of the ellipsoid any more. As we all recognize, the nature is nonlinear and imperfect geometrically. For the purpose of simplification, we usually linearize those system in order to take a step to the future with a good numerical approximation. Actually, many geometrical spherical objects (earth, sun, black hole, etc) are not perfectly sphere in nature. The shape of those objects are closely better interpreted by an ellipsoid because of their rotations by themselves. For example, the ellipsoidal harmonics are represented in calculations of gravitational potential[9]. However spherical harmonic is preferred over the more mathematically complex ellipsoid harmonics (the coefficients in a power series expansions of Lame equation have a relation between three different coefficients).

In 1837, Gabriel Lame introduced second ordinary differential equation which has four regular singular points in the method of separation of variables applied to the Laplace equation in elliptic coordinates[2]. Various authors has called this equation as ‘Lame equation’ or ‘ellipsoidal harmonic equation’[6].

Previously, there was no analytic solution in closed forms of Lame function[6,7,8]. Using Frobenius method to obtain an analytic solution (represented either in the algebraic form or in the Weierstrass’s form), the solution automatically comes out 3 term recurrence relation[7,8]. In contrast, most of well-known special functions consist of two term recursion relation (Hypergeometric, Bessel, Legendre, Kummer functions, etc).

In this paper I’ll construct the power series expansion of Lame function in closed forms analytically and its integral forms with three-term recurrence formula[1]. Lame equation is a second-order linear ordinary differential equation of the algebraic form[2]

$$\frac{\partial^2 y}{\partial x^2}+\frac{1}{2}\left(\frac{1}{x-a}+\frac{1}{x-b}+\frac{1}{x-c}\right)\frac{\partial y}{\partial x}+\frac{-\alpha(\alpha+1)x+q}{4(x-a)(x-b)(x-c)}y=0 \tag{1}$$

Lame’s equation has four regular singular points: a, b, c and ∞. Assume that its solution is

$$y(z)=\sum_{n=0}^{\infty} c_n (x-a)^{n+\lambda} \tag{2}$$

Plug (2) into (1).

$$c_{n+1}=A_n\, c_n+B_n\, c_{n-1} \qquad ;n\geq 1 \tag{3}$$

where,

$$A_n=\frac{\frac{1}{4}(\alpha(\alpha+1)a-q)-(2a-b-c)(n+\lambda)^2}{(a-b)(a-c)(n+1+\lambda)(n+\frac{1}{2}+\lambda)} \tag{4a}$$

$$B_n=\frac{[\alpha-(1-2(n+\lambda))][\alpha-2(n-1+\lambda)]}{2^2(a-b)(a-c)(n+1+\lambda)(n+\frac{1}{2}+\lambda)} \tag{4b}$$

$$c_1=A_0\, c_0 \tag{4c}$$

We have two indicial roots which are $\lambda_1=0$ and $\lambda_2=\frac{1}{2}$

2 Power series

2.1 Polynomial in which makes B_n term terminated

In Ref.[1], the general expression of power series of $y(x)$ for polynomial of x which makes B_n term terminated is

$$
\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} c_n x^{n+\lambda} = \sum_{m=0}^{\infty} y_m(x) = \sum_{l=0}^{\infty} c_l^m x^{l+\lambda} = y_0(x)+y_1(x)+y_2(x)+y_3(x)+\cdots \\
&= c_0 \left\{ \sum_{i_0=0}^{\beta_0} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} \right. \\
&\quad + \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\beta_1} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \right\} x^{2i_2+1+\lambda} \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\beta_0} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\beta_k} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \right) \right. \right. \\
&\quad \left. \left. \left. \times \sum_{i_{2N}=i_{2(N-1)}}^{\beta_N} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \right) \right\} \right\} x^{2i_{2N}+N+\lambda} \right\}
\end{aligned}
\tag{5}
$$

For a polynomial, we need a condition, which is [1]:

$$
B_{2\beta_i+(i+1)} = 0 \qquad \text{where } i, \beta_i = 0,1,2,\cdots \tag{6}
$$

In this paper Pochhammer symbol $(x)_n$ is used to represent the rising factorial: $(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}$. On above, β_i is an eigenvalue that makes B_n term terminated at certain value of n. (6) makes each $y_i(x)$ where $i = 0,1,2,\cdots$ as the polynomial in (5).

2.1.1 The case of $\alpha = 2(2\alpha_i + i + \lambda)$ where $i, \alpha_i = 0,1,2,\cdots$

In (4a)-(4c) replace α by $2(2\alpha_i + i + \lambda)$. In (6) replace index β_i by α_i. Take the new (4a)-(4c), (6) and put them in (5) with replacing variable x by z. After the replacement process, the general expression of power series of $y(z)$ for polynomial in which B_n

term is terminated is

$$
\begin{aligned}
y(z) &= \sum_{m=0}^{\infty} y_m(z) \\
&= c_0 z^{\lambda} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4}+\lambda)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{i_0=0}^{\alpha_0} \left\{ \frac{(2a-b-c)(i_0+\frac{\lambda}{2})^2 - a(\alpha_0+\frac{\lambda}{2})(\alpha_0+\frac{1}{4}+\frac{\lambda}{2}) + \frac{q}{2^4}}{(i_0+\frac{1}{2}+\frac{\lambda}{2})(i_0+\frac{1}{4}+\frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4}+\lambda)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \left. \sum_{i_1=i_0}^{\alpha_1} \left\{ \frac{(-\alpha_1)_{i_1}(\alpha_1+\frac{5}{4}+\lambda)_{i_1}(\frac{3}{2}+\frac{\lambda}{2})_{i_0}(\frac{5}{4}+\frac{\lambda}{2})_{i_0}}{(-\alpha_1)_{i_0}(\alpha_1+\frac{5}{4}+\lambda)_{i_0}(\frac{3}{2}+\frac{\lambda}{2})_{i_1}(\frac{5}{4}+\frac{\lambda}{2})_{i_1}} \eta^{i_1} \right\} \right\} \mu \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(2a-b-c)(i_0+\frac{\lambda}{2})^2 - a(\alpha_0+\frac{\lambda}{2})(\alpha_0+\frac{1}{4}+\frac{\lambda}{2}) + \frac{q}{2^4}}{(i_0+\frac{1}{2}+\frac{\lambda}{2})(i_0+\frac{1}{4}+\frac{\lambda}{2})} \frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4}+\lambda)_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\alpha_k} \frac{(2a-b-c)(i_k+\frac{k}{2}+\frac{\lambda}{2})^2 - a(\alpha_k+\frac{k}{2}+\frac{\lambda}{2})(\alpha_k+\frac{k}{2}+\frac{1}{4}+\frac{\lambda}{2}) + \frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{1}{2}+\frac{\lambda}{2})(i_k+\frac{k}{2}+\frac{1}{4}+\frac{\lambda}{2})} \right. \\
&\quad \times \left. \frac{(-\alpha_k)_{i_k}(\alpha_k+k+\frac{1}{4}+\lambda)_{i_k}(1+\frac{k}{2}+\frac{\lambda}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2}+\frac{\lambda}{2})_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}}(\alpha_k+k+\frac{1}{4}+\lambda)_{i_{k-1}}(1+\frac{k}{2}+\frac{\lambda}{2})_{i_k}(\frac{3}{4}+\frac{k}{2}+\frac{\lambda}{2})_{i_k}} \right\} \\
&\quad \times \left. \left. \sum_{i_n=i_{n-1}}^{\alpha_n} \frac{(-\alpha_n)_{i_n}(\alpha_n+n+\frac{1}{4}+\lambda)_{i_n}(1+\frac{n}{2}+\frac{\lambda}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2}+\frac{\lambda}{2})_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}}(\alpha_n+n+\frac{1}{4}+\lambda)_{i_{n-1}}(1+\frac{n}{2}+\frac{\lambda}{2})_{i_n}(\frac{3}{4}+\frac{n}{2}+\frac{\lambda}{2})_{i_n}} \eta^{i_n} \right\} \mu^n \right\}
\end{aligned}
\tag{7}
$$

where

$$
\begin{cases}
z = x - a \\
\eta = \frac{-(x-a)^2}{(a-b)(a-c)} \\
\mu = \frac{-(x-a)}{(a-b)(a-c)}
\end{cases}
\tag{8}
$$

and

$$
\begin{cases}
\alpha = 2(2\alpha_i + i + \lambda) \ \text{ as } i, \alpha_i = 0,1,2,\cdots \\
\alpha_i \leq \alpha_j \ \text{ only if } i \leq j \ \text{ where } i,j = 0,1,2,\cdots
\end{cases}
\tag{9}
$$

2.1.2 The case of $\alpha = -2(2\alpha_i + i + \lambda) - 1$ *where* $i, \alpha_i = 0,1,2,\cdots$

In (4a)-(4c) replace α by $-2(2\alpha_i + i + \lambda) - 1$. In (6) replace index β_i by α_i. Take the new (4a)-(4c), (6) and put them in (5) with replacing variable x by z. Its solution is exactly equivalent to (7). Put c_0= 1 as λ=0 and $c_0 = \left(\frac{-1}{(a-b)(a-c)} \right)^{\frac{1}{4}}$ as $\lambda = \frac{1}{2}$ in (7). Then, we obtain two independent solutions of Lame equation. The solution is the following ways.

$$\begin{aligned}
y(z) &= LF_{\alpha_j}\left(\alpha_j = \frac{1}{2}\Big(\frac{\alpha}{2}-j\Big)\text{or}-\frac{1}{2}\Big(\frac{\alpha}{2}+\frac{1}{2}+j\Big)\Big|_{j=0,1,2,\cdots};\mu=\frac{-(x-a)}{(a-b)(a-c)};\eta=\frac{-(x-a)^2}{(a-b)(a-c)}\right)\\
&= \sum_{i_0=0}^{\alpha_0}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}}\eta^{i_0} + \sum_{i_0=0}^{\alpha_0}\Bigg\{\frac{(2a-b-c)i_0^2 - a\alpha_0(\alpha_0+\frac{1}{4})+\frac{q}{2^4}}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}}\\
&\quad \times\sum_{i_1=i_0}^{\alpha_1}\Bigg\{\frac{(-\alpha_1)_{i_1}(\alpha_1+\frac{5}{4})_{i_1}(\frac{3}{2})_{i_0}(\frac{5}{4})_{i_0}}{(-\alpha_1)_{i_0}(\alpha_1+\frac{5}{4})_{i_0}(\frac{3}{2})_{i_1}(\frac{5}{4})_{i_1}}\eta^{i_1}\Bigg\}\Bigg\}\mu\\
&\quad +\sum_{n=2}^{\infty}\Bigg\{\sum_{i_0=0}^{\alpha_0}\frac{(2a-b-c)i_0^2 - a\alpha_0(\alpha_0+\frac{1}{4})+\frac{q}{2^4}}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}}\\
&\quad \times\prod_{k=1}^{n-1}\Bigg\{\sum_{i_k=i_{k-1}}^{\alpha_k}\frac{(2a-b-c)(i_k+\frac{k}{2})^2 - a(\alpha_k+\frac{k}{2})(\alpha_k+\frac{k}{2}+\frac{1}{4})+\frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{1}{2})(i_k+\frac{k}{2}+\frac{1}{4})}\\
&\quad \times\frac{(-\alpha_k)_{i_k}(\alpha_k+k+\frac{1}{4})_{i_k}(1+\frac{k}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2})_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}}(\alpha_k+k+\frac{1}{4})_{i_{k-1}}(1+\frac{k}{2})_{i_k}(\frac{3}{4}+\frac{k}{2})_{i_k}}\Bigg\}\\
&\quad \times\sum_{i_n=i_{n-1}}^{\alpha_n}\frac{(-\alpha_n)_{i_n}(\alpha_n+n+\frac{1}{4})_{i_n}(1+\frac{n}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2})_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}}(\alpha_n+n+\frac{1}{4})_{i_{n-1}}(1+\frac{n}{2})_{i_n}(\frac{3}{4}+\frac{n}{2})_{i_n}}\eta^{i_n}\Bigg\}\mu^n
\end{aligned}\tag{10}$$

$$\begin{aligned}
y(z) &= LS_{\alpha_j}\Bigg(\alpha_j=\frac{1}{2}\Big(\frac{\alpha}{2}-\frac{1}{2}-j\Big)\text{or}-\frac{1}{2}\Big(\frac{\alpha}{2}+1+j\Big)\Big|_{j=0,1,2,\cdots};\mu=\frac{-(x-a)}{(a-b)(a-c)}\\
&\quad ;\eta=\frac{-(x-a)^2}{(a-b)(a-c)}\Bigg) = \eta^{\frac{1}{4}}\Bigg\{\sum_{i_0=0}^{\alpha_0}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{3}{4})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}}\eta^{i_0}\\
&\quad +\sum_{i_0=0}^{\alpha_0}\Bigg\{\frac{(2a-b-c)(i_0+\frac{1}{4})^2 - a(\alpha_0+\frac{1}{4})(\alpha_0+\frac{1}{2})+\frac{q}{2^4}}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{3}{4})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}}\\
&\quad \times\sum_{i_1=i_0}^{\alpha_1}\Bigg\{\frac{(-\alpha_1)_{i_1}(\alpha_1+\frac{7}{4})_{i_1}(\frac{7}{4})_{i_0}(\frac{3}{2})_{i_0}}{(-\alpha_1)_{i_0}(\alpha_1+\frac{7}{4})_{i_0}(\frac{7}{4})_{i_1}(\frac{3}{2})_{i_1}}\eta^{i_1}\Bigg\}\Bigg\}\mu\\
&\quad +\sum_{n=2}^{\infty}\Bigg\{\sum_{i_0=0}^{\alpha_0}\frac{(2a-b-c)(i_0+\frac{1}{4})^2 - a(\alpha_0+\frac{1}{4})(\alpha_0+\frac{1}{2})+\frac{q}{2^4}}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{(-\alpha_0)_{i_0}(\alpha_0+\frac{3}{4})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}}\\
&\quad \times\prod_{k=1}^{n-1}\Bigg\{\sum_{i_k=i_{k-1}}^{\alpha_k}\frac{(2a-b-c)(i_k+\frac{k}{2}+\frac{1}{4})^2 - a(\alpha_k+\frac{k}{2}+\frac{1}{4})(\alpha_k+\frac{k}{2}+\frac{1}{2})+\frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{3}{4})(i_k+\frac{k}{2}+\frac{1}{2})}\\
&\quad \times\frac{(-\alpha_k)_{i_k}(\alpha_k+k+\frac{3}{4})_{i_k}(\frac{5}{4}+\frac{k}{2})_{i_{k-1}}(1+\frac{k}{2})_{i_{k-1}}}{(-\alpha_k)_{i_{k-1}}(\alpha_k+k+\frac{3}{4})_{i_{k-1}}(\frac{5}{4}+\frac{k}{2})_{i_k}(1+\frac{k}{2})_{i_k}}\Bigg\}\\
&\quad \times\sum_{i_n=i_{n-1}}^{\alpha_n}\frac{(-\alpha_n)_{i_n}(\alpha_n+n+\frac{3}{4})_{i_n}(\frac{5}{4}+\frac{n}{2})_{i_{n-1}}(1+\frac{n}{2})_{i_{n-1}}}{(-\alpha_n)_{i_{n-1}}(\alpha_n+n+\frac{3}{4})_{i_{n-1}}(\frac{5}{4}+\frac{n}{2})_{i_n}(1+\frac{n}{2})_{i_n}}\eta^{i_n}\Bigg\}\mu^n\Bigg\}
\end{aligned}\tag{11}$$

(10) is the first kind of independent solution of Lame function for the polynomial as $\alpha = 2(2\alpha_j + j)$ or $-2(2\alpha_j + j) - 1$ where $j, \alpha_j = 0, 1, 2, \cdots$. And (11) is the second kind of independent solution of Lame function for the polynomial as $\alpha = 2(2\alpha_j + j) + 1$ or $-2(2\alpha_j + j + 1)$ where $j, \alpha_j = 0, 1, 2, \cdots$.

2.2 Infinite series

In Ref.[1], the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned} y(x) &= \sum_{m=0}^{\infty} y_m(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\ &= c_0 \Bigg\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} + \sum_{i_0=0}^{\infty} \Bigg\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \Bigg\} x^{2i_2+1+\lambda} \\ &\quad + \sum_{N=2}^{\infty} \Bigg\{ \sum_{i_0=0}^{\infty} \Bigg\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \Bigg(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \Bigg) \\ &\quad \times \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \Bigg(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \Bigg) \Bigg\} \Bigg\} x^{2i_{2N}+N+\lambda} \Bigg\} \end{aligned} \tag{12}$$

In (12) replace a variable x by z and substitute (4a)-(4c) into (12). Then, the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned} y(z) &= \sum_{m=0}^{\infty} y_m(z) = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \\ &= c_0 z^{\lambda} \Bigg\{ \sum_{i_0=0}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{\lambda}{2})_{i_0} (\frac{\alpha}{4}+\frac{1}{4}+\frac{\lambda}{2})_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \eta^{i_0} + \sum_{i_0=0}^{\infty} \Bigg\{ \frac{(2a-b-c)(i_0+\frac{\lambda}{2})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{1}{2}+\frac{\lambda}{2})(i_0+\frac{1}{4}+\frac{\lambda}{2})} \\ &\quad \times \frac{(-\frac{\alpha}{4}+\frac{\lambda}{2})_{i_0} (\frac{\alpha}{4}+\frac{1}{4}+\frac{\lambda}{2})_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \sum_{i_1=i_0}^{\infty} \Bigg\{ \frac{(-\frac{\alpha}{4}+\frac{1}{2}+\frac{\lambda}{2})_{i_1} (\frac{\alpha}{4}+\frac{3}{4}+\frac{\lambda}{2})_{i_1} (\frac{3}{2}+\frac{\lambda}{2})_{i_0} (\frac{5}{4}+\frac{\lambda}{2})_{i_0}}{(-\frac{\alpha}{4}+\frac{1}{2}+\frac{\lambda}{2})_{i_0} (\frac{\alpha}{4}+\frac{3}{4}+\frac{\lambda}{2})_{i_0} (\frac{3}{2}+\frac{\lambda}{2})_{i_1} (\frac{5}{4}+\frac{\lambda}{2})_{i_1}} \eta^{i_1} \Bigg\} \Bigg\} \mu \\ &\quad + \sum_{n=2}^{\infty} \Bigg\{ \sum_{i_0=0}^{\infty} \frac{(2a-b-c)(i_0+\frac{\lambda}{2})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{1}{2}+\frac{\lambda}{2})(i_0+\frac{1}{4}+\frac{\lambda}{2})} \frac{(-\frac{\alpha}{4}+\frac{\lambda}{2})_{i_0} (\frac{\alpha}{4}+\frac{1}{4}+\frac{\lambda}{2})_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \\ &\quad \times \prod_{k=1}^{n-1} \Bigg\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(2a-b-c)(i_k+\frac{k}{2}+\frac{\lambda}{2})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{1}{2}+\frac{\lambda}{2})(i_k+\frac{k}{2}+\frac{1}{4}+\frac{\lambda}{2})} \\ &\quad \times \frac{(-\frac{\alpha}{4}+\frac{k}{2}+\frac{\lambda}{2})_{i_k} (\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{4}+\frac{\lambda}{2})_{i_k} (1+\frac{k}{2}+\frac{\lambda}{2})_{i_{k-1}} (\frac{k}{2}+\frac{3}{4}+\frac{\lambda}{2})_{i_{k-1}}}{(-\frac{\alpha}{4}+\frac{k}{2}+\frac{\lambda}{2})_{i_{k-1}} (\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{4}+\frac{\lambda}{2})_{i_{k-1}} (1+\frac{k}{2}+\frac{\lambda}{2})_{i_k} (\frac{k}{2}+\frac{3}{4}+\frac{\lambda}{2})_{i_k}} \Bigg\} \\ &\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{n}{2}+\frac{\lambda}{2})_{i_n} (\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{4}+\frac{\lambda}{2})_{i_n} (1+\frac{n}{2}+\frac{\lambda}{2})_{i_{n-1}} (\frac{n}{2}+\frac{3}{4}+\frac{\lambda}{2})_{i_{n-1}}}{(-\frac{\alpha}{4}+\frac{n}{2}+\frac{\lambda}{2})_{i_{n-1}} (\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{4}+\frac{\lambda}{2})_{i_{n-1}} (1+\frac{n}{2}+\frac{\lambda}{2})_{i_n} (\frac{n}{2}+\frac{3}{4}+\frac{\lambda}{2})_{i_n}} \eta^{i_n} \Bigg\} \mu^n \Bigg\} \end{aligned} \tag{13}$$

Put c_0= 1 as λ=0 and $c_0 = \left(\frac{-1}{(a-b)(a-c)} \right)^{\frac{1}{4}}$ as $\lambda = \frac{1}{2}$ in (13). Then, we obtain two independent solutions of Lame equation. The solution is the following ways.

$$\begin{aligned}
y(z) &= LF\left(\mu = \frac{-(x-a)}{(a-b)(a-c)}; \eta = \frac{-(x-a)^2}{(a-b)(a-c)}\right) \\
&= \sum_{i_0=0}^{\infty} \frac{(-\frac{\alpha}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}} \eta^{i_0} + \sum_{i_0=0}^{\infty} \left\{ \frac{(2a-b-c)i_0^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})} \right. \\
&\quad \left. \times \frac{(-\frac{\alpha}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}} \sum_{i_1=i_0}^{\infty} \left\{ \frac{(-\frac{\alpha}{4}+\frac{1}{2})_{i_1}(\frac{\alpha}{4}+\frac{3}{4})_{i_1}(\frac{3}{2})_{i_0}(\frac{5}{4})_{i_0}}{(-\frac{\alpha}{4}+\frac{1}{2})_{i_0}(\frac{\alpha}{4}+\frac{3}{4})_{i_0}(\frac{3}{2})_{i_1}(\frac{5}{4})_{i_1}} \eta^{i_1} \right\} \right\} \mu \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(2a-b-c)i_0^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})} \frac{(-\frac{\alpha}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{4})_{i_0}}{(\frac{3}{4})_{i_0}(1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(2a-b-c)(i_k+\frac{k}{2})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{1}{2})(i_k+\frac{k}{2}+\frac{1}{4})} \right. \\
&\quad \left. \times \frac{(-\frac{\alpha}{4}+\frac{k}{2})_{i_k}(\frac{\alpha}{4}+\frac{1}{4}+\frac{k}{2})_{i_k}(1+\frac{k}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2})_{i_{k-1}}}{(-\frac{\alpha}{4}+\frac{k}{2})_{i_{k-1}}(\frac{\alpha}{4}+\frac{1}{4}+\frac{k}{2})_{i_{k-1}}(1+\frac{k}{2})_{i_k}(\frac{3}{4}+\frac{k}{2})_{i_k}} \right\} \\
&\quad \left. \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{n}{2})_{i_n}(\frac{\alpha}{4}+\frac{1}{4}+\frac{n}{2})_{i_n}(1+\frac{n}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2})_{i_{n-1}}}{(-\frac{\alpha}{4}+\frac{n}{2})_{i_{n-1}}(\frac{\alpha}{4}+\frac{1}{4}+\frac{n}{2})_{i_{n-1}}(1+\frac{n}{2})_{i_n}(\frac{3}{4}+\frac{n}{2})_{i_n}} \eta^{i_n} \right\} \mu^n
\end{aligned} \tag{14}$$

$$\begin{aligned}
y(z) &= LS\left(\mu = \frac{-(x-a)}{(a-b)(a-c)}; \eta = \frac{-(x-a)^2}{(a-b)(a-c)}\right) \\
&= \eta^{\frac{1}{4}} \left\{ \sum_{i_0=0}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{1}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{2})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}} \eta^{i_0} + \sum_{i_0=0}^{\infty} \left\{ \frac{(2a-b-c)(i_0+\frac{1}{4})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})} \right. \right. \\
&\quad \left. \times \frac{(-\frac{\alpha}{4}+\frac{1}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{2})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}} \sum_{i_1=i_0}^{\infty} \left\{ \frac{(-\frac{\alpha}{4}+\frac{3}{4})_{i_1}(\frac{\alpha}{4}+1)_{i_1}(\frac{7}{4})_{i_0}(\frac{3}{2})_{i_0}}{(-\frac{\alpha}{4}+\frac{3}{4})_{i_0}(\frac{\alpha}{4}+1)_{i_0}(\frac{7}{4})_{i_1}(\frac{3}{2})_{i_1}} \eta^{i_1} \right\} \right\} \mu \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \frac{(2a-b-c)(i_0+\frac{1}{4})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})} \frac{(-\frac{\alpha}{4}+\frac{1}{4})_{i_0}(\frac{\alpha}{4}+\frac{1}{2})_{i_0}}{(\frac{5}{4})_{i_0}(1)_{i_0}} \right. \\
&\quad \times \prod_{k=1}^{n-1} \left\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(2a-b-c)(i_k+\frac{k}{2}+\frac{1}{4})^2 - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4}}{(i_k+\frac{k}{2}+\frac{3}{4})(i_k+\frac{k}{2}+\frac{1}{2})} \right. \\
&\quad \left. \times \frac{(-\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{4})_{i_k}(\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{2})_{i_k}(\frac{5}{4}+\frac{k}{2})_{i_{k-1}}(1+\frac{k}{2})_{i_{k-1}}}{(-\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{4})_{i_{k-1}}(\frac{\alpha}{4}+\frac{k}{2}+\frac{1}{2})_{i_{k-1}}(\frac{5}{4}+\frac{k}{2})_{i_k}(1+\frac{k}{2})_{i_k}} \right\} \\
&\quad \left. \left. \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{4})_{i_n}(\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{2})_{i_n}(\frac{5}{4}+\frac{n}{2})_{i_{n-1}}(1+\frac{n}{2})_{i_{n-1}}}{(-\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{4})_{i_{n-1}}(\frac{\alpha}{4}+\frac{n}{2}+\frac{1}{2})_{i_{n-1}}(\frac{5}{4}+\frac{n}{2})_{i_n}(1+\frac{n}{2})_{i_n}} \eta^{i_n} \right\} \mu^n \right\}
\end{aligned} \tag{15}$$

(14) is the first kind of independent solution of Lame function for the infinite series. And (15) is the second kind of independent solution of Lame function for the infinite series.

3 Asymptotic behavior of the function $y(z = x - a)$ and the boundary condition for x

3.1 Infinite series

Now let's test for convergence of infinite series of the analytic function $y(z)$. As n goes to infinity, (4a) and (4b) are

$$\lim_{n\gg 1} A_n = A = \frac{-(2a-b-c)}{(a-b)(a-c)} \qquad \lim_{n\gg 1} B_n = B = \frac{-1}{(a-b)(a-c)} \tag{16}$$

Substitute (16) into (3). For $n = 0, 1, 2, \cdots$, it give

$$\begin{aligned}
& c_0 \\
& c_1 = AC_0 \\
& c_2 = (A^2 + B)C_0 \\
& c_3 = (A^3 + 2AB)C_0 \\
& c_4 = (A^4 + 3A^2B + B^2)C_0 \\
& c_5 = (A^5 + 4A^3B + 3AB^2)C_0 \\
& c_6 = (A^6 + 5A^4B + 6A^2B^2 + B^3)C_0 \\
& c_7 = (A^7 + 6A^5B + 10A^3B^2 + 4AB^3)C_0 \\
& c_8 = (A^8 + 7A^6B + 15A^4B^2 + 10A^2B^3 + B^4)C_0 \\
& \quad \vdots \qquad\qquad\qquad \vdots
\end{aligned} \tag{17}$$

The sequences c_n consists of combinations A and B in (17). First observe the term inside parentheses of sequence c_n which does not include any A_n's: c_n with even index (c_0,c_2, c_4,$\cdots$).

(1) Zero term of A's

$$\begin{aligned}
& c_0 \\
& c_2 = Bc_0 \\
& c_4 = B^2c_0 \\
& c_6 = B^3c_0 \\
& c_8 = B^4c_0 \\
& c_{10} = B^5c_0 \\
& \;\vdots \qquad \vdots
\end{aligned} \tag{18}$$

When a function $y(z)$, analytic at z=0, is expanded in a power series z=0, we write

$$y(z)=\sum_{m=0}^{\infty} y_m(z) \tag{19}$$

where

$$y_m(z)=\sum_{n=0}^{\infty} c_n^m z^n \tag{20}$$

Put(18) in (20) putting $m=0$.

$$y_0(z)=c_0\sum_{n=0}^{\infty}\left(Bz^2\right)^n \tag{21}$$

Observe the terms inside parentheses of sequence c_n which include one term of A_n's in (17): c_n with odd index (c_1, c_3, c_5,$\cdots$).

(2) One term of A's

$$\begin{aligned} c_1 &= Ac_0 \\ c_3 &= 2ABc_0 \\ c_5 &= 3AB^2c_0 \\ c_7 &= 4AB^3c_0 \\ c_9 &= 5AB^4c_0 \\ \vdots &\qquad \vdots \end{aligned} \tag{22}$$

Put (22) in (20) putting $m=1$.

$$y_1(z)=c_0Az\sum_{n=0}^{\infty}\frac{(n+1)}{1!}\left(Bz^2\right)^n \tag{23}$$

Observe the terms inside parentheses of sequence c_n which include two terms of A_n's in (17): c_n with even index (c_2, c_4, c_6,$\cdots$).

(3) Two terms of A's

$$\begin{aligned} c_2 &= A^2c_0 \\ c_4 &= 3A^2Bc_0 \\ c_6 &= 6A^2B^2c_0 \\ c_8 &= 10A^2B^3c_0 \\ c_{10} &= 15A^2B^4c_0 \\ \vdots &\qquad \vdots \end{aligned} \tag{24}$$

Put (23) in (20) putting $m=2$.

$$y_2(z)=c_0\left(Az\right)^2\sum_{n=0}^{\infty}\frac{(n+1)(n+2)}{2!}\left(Bz^2\right)^n \tag{25}$$

By repeating this process for all higher terms of A's, I obtain every $y_m(x)$ terms where $m \geq 3$. Substitute (21), (23), (25) and including all $y_m(x)$ terms where $m \geq 3$ into (19).

$$\begin{aligned} y(z) &= \sum_{n=0}^{\infty} c_n z^{n+\lambda} = y_0(z) + y_1(z) + y_2(z) + y_3(z) + \cdots \qquad (26) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(n+m)!}{n!\, m!} \tilde{x}^n \tilde{y}^m = \frac{1}{1-(\tilde{x}+\tilde{y})} \qquad \text{where } c_0 = 1, \tilde{x} = Bz^2 \text{ and } \tilde{y} = Az \end{aligned}$$

Substitute (16) in (26).

$$\lim_{n \gg 1} y(z) = \frac{1}{1 + \left(\frac{(x-a)^2}{(a-b)(a-c)} + \frac{(2a-b-c)(x-a)}{(a-b)(a-c)} \right)} \qquad (27)$$

(27) is geometric series. The condition of convergence of it is

$$\left| \frac{(x-a)^2}{(a-b)(a-c)} + \frac{(2a-b-c)(x-a)}{(a-b)(a-c)} \right| < 1 \qquad (28)$$

According to (28), we have a boundary condition of x for the infinite series of Lame function in which is following the way.

(A) As a=b or a=c

$$\text{no solution} \qquad (29a)$$

(B) As $0 < \left(a - \frac{b+c}{2}\right)^2 < (a-b)(a-c)$

$$\frac{(b+c)}{2} - \sqrt{\left(a - \frac{b+c}{2}\right)^2 + (a-b)(a-c)} < x < \frac{(b+c)}{2} + \sqrt{\left(a - \frac{b+c}{2}\right)^2 + (a-b)(a-c)} \qquad (29b)$$

(C) As $(a-b)(a-c) = \left(a - \frac{b+c}{2}\right)^2 > 0$

$$\frac{(b+c)}{2} - \sqrt{2(a-b)(a-c)} < x < \frac{(b+c)}{2} \quad \text{and} \quad \frac{(b+c)}{2} < x < \frac{(b+c)}{2} + \sqrt{2(a-b)(a-c)} \qquad (29c)$$

(D) As $0 < (a-b)(a-c) < \left(a - \frac{b+c}{2}\right)^2$

$$\begin{cases} \frac{(b+c)}{2} - \sqrt{\left(a - \frac{b+c}{2}\right)^2 + (a-b)(a-c)} < x < \frac{(b+c)}{2} - \sqrt{\left(a - \frac{b+c}{2}\right)^2 - (a-b)(a-c)} \\ \frac{(b+c)}{2} + \sqrt{\left(a - \frac{b+c}{2}\right)^2 - (a-b)(a-c)} < x < \frac{(b+c)}{2} + \sqrt{\left(a - \frac{b+c}{2}\right)^2 + (a-b)(a-c)} \end{cases} \qquad (29d)$$

(E) As $0 < \left(a - \frac{b+c}{2}\right)^2 < -(a-b)(a-c)$

$$\frac{(b+c)}{2} - \sqrt{\left(a - \frac{b+c}{2}\right)^2 - (a-b)(a-c)} < x < \frac{(b+c)}{2} + \sqrt{\left(a - \frac{b+c}{2}\right)^2 - (a-b)(a-c)} \qquad (29e)$$

(F) As $-(a-b)(a-c)=\left(a-\frac{b+c}{2}\right)^2>0$

$$\frac{(b+c)}{2}-\sqrt{-2(a-b)(a-c)}<x<\frac{(b+c)}{2}\quad\text{and}\quad\frac{(b+c)}{2}<x<\frac{(b+c)}{2}+\sqrt{-2(a-b)(a-c)} \tag{29f}$$

(G) As $0<-(a-b)(a-c)<\left(a-\frac{b+c}{2}\right)^2$

$$\begin{cases}\frac{(b+c)}{2}-\sqrt{\left(a-\frac{b+c}{2}\right)^2-(a-b)(a-c)}<x<\frac{(b+c)}{2}-\sqrt{\left(a-\frac{b+c}{2}\right)^2+(a-b)(a-c)}\\ \frac{(b+c)}{2}+\sqrt{\left(a-\frac{b+c}{2}\right)^2+(a-b)(a-c)}<x<\frac{(b+c)}{2}+\sqrt{\left(a-\frac{b+c}{2}\right)^2-(a-b)(a-c)}\end{cases} \tag{29g}$$

3.2 The case of $2a-b-c\gg 1$ or $2a-b-c\ll -1$

Let assume that $2a-b-c$ is much greater than 1 or is much less than -1. Then B_n terms are negligible. (3) is approximately

$$c_{n+1}\approx A_n c_n \tag{30a}$$

And,

$$\lim_{n\gg 1} A_n\approx A=\frac{-(2a-b-c)}{(a-b)(a-c)}\qquad c_1=A_0c_0\approx Ac_0 \tag{30b}$$

Substitute (30b) into (30a). For $n=0,1,2,\cdots$, it give

$$\begin{aligned}&c_0\\ &c_1=AC_0\\ &c_2=A^2C_0\\ &c_3=A^3C_0\\ &c_4=A^4C_0\\ &\vdots\qquad\vdots\end{aligned} \tag{31}$$

When a function $y(z)$, analytic at z=0, is expanded in a power series z=0 by using (31), we write

$$\lim_{n\gg 1} y(z)=c_0\sum_{n=0}^{\infty}(Az)^n=\frac{1}{1-Az}\qquad\text{where } |Az|<1 \text{ and } c_0=1 \tag{32}$$

Substitute (30b) into (32)

$$\lim_{n\gg 1} y(z)=\frac{1}{1+\left(\frac{(2a-b-c)(x-a)}{(a-b)(a-c)}\right)}\qquad\text{where } a\neq b \text{ and } a\neq c \tag{33}$$

(33) is binomial series. The condition of convergence of x is

$$\left|\frac{(2a-b-c)(x-a)}{(a-b)(a-c)}\right| < 1 \tag{34}$$

3.3 The case of $2a-b-c \approx 0$

Let assume that $2a-b-c$ is approximately close to 0. But $a \neq 0$ in (5). Then A_n terms are negligible. (3) is approximately

$$c_{n+1} \approx B_n c_{n-1} \qquad \text{where } n \geq 1 \tag{35a}$$

And,

$$\lim_{n\gg 1} B_n \approx B = \frac{-1}{(a-b)(a-c)} \tag{35b}$$

We can classify c_n as to even and odd terms from plugging (35b) into (35a).

$$\begin{array}{ll} c_0 & c_1 \\ c_2 = Bc_0 & c_3 = Bc_1 \\ c_4 = B^2c_0 & c_5 = B^2c_1 \\ c_6 = B^3c_0 & c_7 = B^3c_1 \\ c_8 = B^4c_0 & c_9 = B^4c_1 \\ c_{10} = B^5c_0 & c_{11} = B^5c_1 \\ \vdots & \vdots \end{array} \tag{36}$$

When a function $y(z)$, analytic at z=0, is expanded in a power series z=0 by using (36), we write

$$\lim_{n\gg 1} y(z) = c_0 \sum_{n=0}^{\infty} \left(Bz^2\right)^n + c_1 z \sum_{n=0}^{\infty} \left(Bz^2\right)^n = c_0 \frac{1}{1-Bz^2} + c_1 \frac{z}{1-Bz^2} \qquad \text{where } |Bz^2| < 1 \tag{37}$$

Substitute (35b) into (37). And for simplicity, let say $c_0 = c_1 = 1$ into it.

$$\lim_{n\gg 1} y(z) = \frac{1+(x-a)}{1+\left(\frac{(x-a)^2}{(a-b)(a-c)}\right)} \qquad \text{where } a \neq b \text{ and } a \neq c \tag{38}$$

The condition of convergence of x is

$$\left|\frac{(x-a)^2}{(a-b)(a-c)}\right| < 1 \tag{39}$$

4 Integral Formalism

4.1 Polynomial in which makes B_n term terminated

Now let's investigate the integral formalism for the polynomial case of B_n term terminated at certain eigenvalue. There is a generalized hypergeometric function which is: In this paper Pochhammer symbol $(x)_n$ is used to represent the rising factorial: $(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}$.

$$\begin{aligned} I_l &= \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l}(\alpha_l+l+\frac{1}{4}+\lambda)_{i_l}(1+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}}(\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}}(\alpha_l+l+\frac{1}{4}+\lambda)_{i_{l-1}}(1+\frac{l}{2}+\frac{\lambda}{2})_{i_l}(\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_l}}\eta^{i_l} \\ &= \eta^{i_{l-1}} \sum_{j=0}^{\infty} \frac{B(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2},j+1)B(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2},j+1)(i_{l-1}-\alpha_l)_j(\alpha_l+i_{l-1}+l+\frac{1}{4}+\lambda)_j}{(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2})^{-1}(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2})^{-1}(1)_j\; j!}\eta^j \end{aligned} \tag{40}$$

By using integral form of beta function,

$$B\left(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2},j+1\right) = \int_0^1 dt_l\, t_l^{i_{l-1}+\frac{l}{2}-\frac{5}{4}+\frac{\lambda}{2}}(1-t_l)^j \tag{41a}$$

$$B\left(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2},j+1\right) = \int_0^1 du_l\, u_l^{i_{l-1}+\frac{l}{2}-1+\frac{\lambda}{2}}(1-u_l)^j \tag{41b}$$

Substitute (41a) and (41b) into (40). And divide $(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2})(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2})$ into I_l.

$$\begin{aligned} &\frac{1}{(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2})(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2})} \\ &\quad\times \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l}(\alpha_l+l+\frac{1}{4}+\lambda)_{i_l}(1+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}}(\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}}(\alpha_l+l+\frac{1}{4}+\lambda)_{i_{l-1}}(1+\frac{l}{2}+\frac{\lambda}{2})_{i_l}(\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_l}}\eta^{i_l} \\ &= \int_0^1 dt_l\, t_l^{\frac{l}{2}-\frac{5}{4}+\frac{\lambda}{2}} \int_0^1 du_l\, u_l^{\frac{l}{2}-1+\frac{\lambda}{2}} (t_l u_l \eta)^{i_{l-1}} \\ &\quad\times \sum_{j=0}^{\infty} \frac{(i_{l-1}-\alpha_l)_j(i_{l-1}+l+\alpha_l+\frac{1}{4}+\lambda)_j}{(1)_j\; j!}[\eta(1-t_l)(1-u_l)]^j \end{aligned} \tag{42}$$

The integral form of hypergeometric function is

$$\begin{aligned} {}_2F_1(\alpha,\beta;\gamma;z) &= \sum_{n=0}^{\infty} \frac{(\alpha)_n(\beta)_n}{(\gamma)_n(n!)}z^n \\ &= -\frac{1}{2\pi i}\frac{\Gamma(1-\alpha)\Gamma(\gamma)}{\Gamma(\gamma-\alpha)}\oint dv_l\,(-v_l)^{\alpha-1}(1-v_l)^{\gamma-\alpha-1}(1-zv_l)^{-\beta} \\ &\quad \text{where } \mathrm{Re}(\gamma-\alpha)>0 \end{aligned} \tag{43}$$

replaced α, β, γ and z by $i_{l-1}-\alpha_l$, $i_{l-1}+l+\alpha_l+\frac{l}{4}+\lambda$, 1 and $\eta(1-t_l)(1-u_l)$ in (43)

$$\begin{aligned}
&\sum_{j=0}^{\infty} \frac{(i_{l-1}-\alpha_l)_j (i_{l-1}+l+\alpha_l+\frac{l}{4}+\lambda)_j}{(1)_j \; j!} [\eta(1-t_l)(1-u_l)]^j \\
&= \frac{1}{2\pi i} \oint dv_l \; \frac{1}{v_l} (1-\eta v_l(1-t_l)(1-u_l))^{-(l+\frac{1}{4}+\lambda)} \left(\frac{1-\frac{1}{v_l}}{1-\eta v_l(1-t_l)(1-u_l)} \right)^{\alpha_l} \\
&\times \left(\frac{1}{(1-\frac{1}{v_l})(1-\eta v_l(1-t_l)(1-u_l))} \right)^{i_{l-1}} \qquad (44)
\end{aligned}$$

Substitute (44) into (42).

$$\begin{aligned}
&\frac{1}{(i_{l-1}+\frac{l}{2}-\frac{1}{4}+\frac{\lambda}{2})(i_{l-1}+\frac{l}{2}+\frac{\lambda}{2})} \\
&\times \sum_{i_l=i_{l-1}}^{\alpha_l} \frac{(-\alpha_l)_{i_l} (\alpha_l+l+\frac{1}{4}+\lambda)_{i_l} (1+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}} (\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_{l-1}}}{(-\alpha_l)_{i_{l-1}} (\alpha_l+l+\frac{1}{4}+\lambda)_{i_{l-1}} (1+\frac{l}{2}+\frac{\lambda}{2})_{i_l} (\frac{3}{4}+\frac{l}{2}+\frac{\lambda}{2})_{i_l}} \eta^{i_l} \\
&= \int_0^1 dt_l \; t_l^{\frac{l}{2}-\frac{5}{4}+\frac{\lambda}{2}} \int_0^1 du_l \; u_l^{\frac{l}{2}-1+\frac{\lambda}{2}} \frac{1}{2\pi i} \oint dv_l \; \frac{1}{v_l} (1-\eta v_l(1-t_l)(1-u_l))^{-(l+\frac{1}{4}+\lambda)} \\
&\times \left(\frac{1-\frac{1}{v_l}}{1-\eta v_l(1-t_l)(1-u_l)} \right)^{\alpha_l} \left(\frac{t_l u_l v_l}{(v_l-1)} \frac{\eta}{1-\eta v_l(1-t_l)(1-u_l)} \right)^{i_{l-1}} \qquad (45)
\end{aligned}$$

Substitute (45) into (7) where $l = 1,2,3,\cdots$, and the integral formalism of Lame function for polynomial in which B_n term terminated is

$$\begin{aligned}
y(z) &= \sum_{m=0}^{\infty} y_m(z) \\
&= c_0 z^{\lambda} \left\{ \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0} (\alpha_0+\frac{1}{4}+\lambda)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \eta^{i_0} \right. \\
&\quad + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left(\int_0^1 dt_{n-k} \; t_{n-k}^{\frac{1}{2}(n-k-\frac{5}{2}+\lambda)} \int_0^1 du_{n-k} \; u_{n-k}^{\frac{1}{2}(n-k-2+\lambda)} \right. \right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})\right)^{-(n-k+\frac{1}{4}+\lambda)} \\
&\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})} \right)^{\alpha_{n-k}} \\
&\quad \times \left\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-1+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-1+\lambda)} \right. \\
&\quad \left. \left. -a \left(\alpha_{n-k-1}+\frac{1}{2}(n-k-1+\lambda) \right) \left(\alpha_{n-k-1}+\frac{1}{2}(n-k-\frac{1}{2}+\lambda) \right) + \frac{q}{2^4} \right\} \right) \\
&\quad \left. \left. \times \sum_{i_0=0}^{\alpha_0} \frac{(-\alpha_0)_{i_0} (\alpha_0+\frac{1}{4}+\lambda)_{i_0}}{(1+\frac{\lambda}{2})_{i_0} (\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \right\} \mu^n \right\} \qquad (46)
\end{aligned}$$

where

$$\overleftrightarrow{w}_{i,j} = \begin{cases} \dfrac{1}{(v_i - 1)} \dfrac{\overleftrightarrow{w}_{i+1,j} v_i t_i u_i}{1 - \overleftrightarrow{w}_{i+1,j} v_i (1-t_i)(1-u_i)} \\ \eta \ \text{ only if } i > j \end{cases} \tag{47}$$

Put c_0= 1 as λ=0 and $c_0 = \left(\dfrac{-1}{(a-b)(a-c)} \right)^{\frac{1}{4}}$ as $\lambda = \frac{1}{2}$ in (46). Also apply (43) into it. Then I obtain two independent solutions of Lame equation. The solution is the following ways.

$$\begin{aligned} y(z) &= LF_{\alpha_j} \left(\alpha_j = \frac{1}{2}\Big(\frac{\alpha}{2} - j\Big) \text{or} - \frac{1}{2}\Big(\frac{\alpha}{2} + \frac{1}{2} + j\Big)\Big|_{j=0,1,2,\cdots} ; \mu = \frac{-(x-a)}{(a-b)(a-c)} \right. \\ &\qquad \left. ; \eta = \frac{-(x-a)^2}{(a-b)(a-c)} \right) \\ &= {}_2F_1 \left(-\alpha_0, \alpha_0 + \frac{1}{4}; \frac{3}{4}; \eta \right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left(\int_0^1 dt_{n-k}\, t_{n-k}^{\frac{1}{2}(n-k-\frac{5}{2})} \int_0^1 du_{n-k}\, u_{n-k}^{\frac{1}{2}(n-k-2)} \right. \right. \\ &\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1 - \overleftrightarrow{w}_{n-k+1,n} v_{n-k} (1-t_{n-k})(1-u_{n-k}) \right)^{-(n-k+\frac{1}{4})} \\ &\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1 - \overleftrightarrow{w}_{n-k+1,n} v_{n-k} (1-t_{n-k})(1-u_{n-k})} \right)^{\alpha_{n-k}} \\ &\quad \times \Bigg\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-1)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-1)} \\ &\quad \left. -a \left(\alpha_{n-k-1} + \frac{1}{2}(n-k-1) \right) \left(\alpha_{n-k-1} + \frac{1}{2}\Big(n-k-\frac{1}{2}\Big) \right) + \frac{q}{2^4} \Bigg\} \right) \\ &\quad \left. \times {}_2F_1 \left(-\alpha_0, \alpha_0 + \frac{1}{4}; \frac{3}{4}; \overleftrightarrow{w}_{1,n} \right) \right\} \mu^n \end{aligned} \tag{48}$$

$$
\begin{aligned}
y(z) &= LS_{\alpha_j}\Bigg(\alpha_j = \frac{1}{2}\Big(\frac{\alpha}{2}-\frac{1}{2}-j\Big)\text{or}-\frac{1}{2}\Big(\frac{\alpha}{2}+1+j\Big)\Big|_{j=0,1,2,\cdots} ;\mu = \frac{-(x-a)}{(a-b)(a-c)} \\
&\qquad ;\eta = \frac{-(x-a)^2}{(a-b)(a-c)}\Bigg) \\
&= \eta^{\frac{1}{4}} \Bigg\{ {}_2F_1\left(-\alpha_0, \alpha_0+\frac{3}{4};\frac{5}{4};\eta\right) \\
&\quad + \sum_{n=1}^{\infty} \Bigg\{ \prod_{k=0}^{n-1} \Bigg(\int_0^1 dt_{n-k}\, t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k}\, u_{n-k}^{\frac{1}{2}(n-k-\frac{3}{2})} \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})\right)^{-(n-k+\frac{3}{4})} \\
&\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})} \right)^{\alpha_{n-k}} \\
&\quad \times \Bigg\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-\frac{1}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-\frac{1}{2})} \\
&\quad -a\left(\alpha_{n-k-1}+\frac{1}{2}\Big(n-k-\frac{1}{2}\Big)\right) \left(\alpha_{n-k-1}+\frac{1}{2}(n-k)\right) + \frac{q}{2^4} \Bigg\} \Bigg) \\
&\quad \times {}_2F_1\left(-\alpha_0, \alpha_0+\frac{3}{4};\frac{5}{4};\overleftrightarrow{w}_{1,n}\right) \Bigg\} \mu^n \Bigg\}
\end{aligned}
\tag{49}
$$

(48) is the integral formalism of the first kind of independent solution of Lame function for the polynomial as $\alpha = 2(2\alpha_j+j)$ or $-2(2\alpha_j+j)-1$ where $j, \alpha_j = 0,1,2,\cdots$. And, (49) is the integral formalism of the second kind of independent solution of Lame function for the polynomial as $\alpha = 2(2\alpha_j+j)+1$ or $-2(2\alpha_j+j+1)$ where $j, \alpha_j = 0,1,2,\cdots$.

4.2 Infinite series

For infinite series, replace the finite summation with an interval $[0,\alpha_0]$ by infinite summation with an interval $[0,\infty]$ in (46). Also, replaced α_0, α_{n-k} and α_{n-k-1} by

$\frac{1}{2}\left(\frac{\alpha}{2}-\lambda\right)$, $\frac{1}{2}\left(\frac{\alpha}{2}-n+k-\lambda\right)$ and $\frac{1}{2}\left(\frac{\alpha}{2}-n+k+1-\lambda\right)$ into (46).

$$\begin{aligned}
y(z) &= \sum_{n=0}^{\infty} y_n(z) \\
&= c_0 z^{\lambda} \Bigg\{ \sum_{i_0=0}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{\lambda}{2})_{i_0}(\frac{\alpha}{4}+\frac{1}{4}+\frac{\lambda}{2})_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \eta^{i_0} \\
&\quad + \sum_{n=1}^{\infty} \Bigg\{ \prod_{k=0}^{n-1} \Bigg(\int_0^1 dt_{n-k}\, t_{n-k}^{\frac{1}{2}(n-k-\frac{5}{2}+\lambda)} \int_0^1 du_{n-k}\, u_{n-k}^{\frac{1}{2}(n-k-2+\lambda)} \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})\right)^{-(n-k+\frac{1}{4}+\lambda)} \\
&\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})} \right)^{\frac{1}{2}(\frac{\alpha}{2}-n+k-\lambda)} \\
&\quad \times \Bigg\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-1+\lambda)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-1+\lambda)} \\
&\quad - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4} \Bigg\} \Bigg) \sum_{i_0=0}^{\infty} \frac{(-\frac{\alpha}{4}+\frac{\lambda}{2})_{i_0}(\frac{\alpha}{4}+\frac{1}{4}+\frac{\lambda}{2})_{i_0}}{(1+\frac{\lambda}{2})_{i_0}(\frac{3}{4}+\frac{\lambda}{2})_{i_0}} \overleftrightarrow{w}_{1,n}^{i_0} \Bigg\} \mu^n \Bigg\}
\end{aligned} \tag{50}$$

Put c_0= 1 as λ=0 and $c_0 = \left(\frac{-1}{(a-b)(a-c)}\right)^{\frac{1}{4}}$ as $\lambda = \frac{1}{2}$ in eq(118). Also apply eq(99) into it. Then I obtain two independent solutions of Lame equation. The solution is the following ways.

$$\begin{aligned}
y(z) &= LF\left(\mu = \frac{-(x-a)}{(a-b)(a-c)}; \eta = \frac{-(x-a)^2}{(a-b)(a-c)} \right) \\
&= {}_2F_1\left(-\frac{\alpha}{4}, \frac{\alpha}{4}+\frac{1}{4}; \frac{3}{4}; \eta\right) + \sum_{n=1}^{\infty} \Bigg\{ \prod_{k=0}^{n-1} \Bigg(\int_0^1 dt_{n-k}\, t_{n-k}^{\frac{1}{2}(n-k-\frac{5}{2})} \int_0^1 du_{n-k}\, u_{n-k}^{\frac{1}{2}(n-k-2)} \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})\right)^{-(n-k+\frac{1}{4})} \\
&\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})} \right)^{\frac{1}{2}(\frac{\alpha}{2}-n+k)} \\
&\quad \times \Bigg\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-1)} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-1)} - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4} \Bigg\} \Bigg) \\
&\quad \times {}_2F_1\left(-\frac{\alpha}{4}, \frac{\alpha}{4}+\frac{1}{4}; \frac{3}{4}; \overleftrightarrow{w}_{1,n}\right) \Bigg\} \mu^n
\end{aligned} \tag{51}$$

$$
\begin{aligned}
y(z) &= LS\left(\mu = \frac{-(x-a)}{(a-b)(a-c)}; \eta = \frac{-(x-a)^2}{(a-b)(a-c)}\right) \\
&= \eta^{\frac{1}{4}} \left\{ {}_2F_1\left(-\frac{\alpha}{4}+\frac{1}{4}, \frac{\alpha}{4}+\frac{1}{2}; \frac{5}{4}; \eta\right) + \sum_{n=1}^{\infty} \left\{ \prod_{k=0}^{n-1} \left(\int_0^1 dt_{n-k}\, t_{n-k}^{\frac{1}{2}(n-k-2)} \int_0^1 du_{n-k}\, u_{n-k}^{\frac{1}{2}(n-k-\frac{3}{2})} \right.\right.\right. \\
&\quad \times \frac{1}{2\pi i} \oint dv_{n-k} \frac{1}{v_{n-k}} \left(1 - \overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})\right)^{-(n-k+\frac{3}{4})} \\
&\quad \times \left(\frac{(v_{n-k}-1)}{v_{n-k}} \frac{1}{1-\overleftrightarrow{w}_{n-k+1,n} v_{n-k}(1-t_{n-k})(1-u_{n-k})} \right)^{\frac{1}{2}(\frac{\alpha}{2}-n+k-\frac{1}{2})} \\
&\quad \left.\times \left\{ (2a-b-c) \overleftrightarrow{w}_{n-k,n}^{-\frac{1}{2}(n-k-\frac{1}{2})} \left(\overleftrightarrow{w}_{n-k,n} \partial_{\overleftrightarrow{w}_{n-k,n}} \right)^2 \overleftrightarrow{w}_{n-k,n}^{\frac{1}{2}(n-k-\frac{1}{2})} - \frac{a}{2^4}\alpha(\alpha+1) + \frac{q}{2^4} \right\} \right) \\
&\quad \left.\left.\times {}_2F_1\left(-\frac{\alpha}{4}+\frac{1}{4}, \frac{\alpha}{4}+\frac{1}{2}; \frac{5}{4}; \overleftrightarrow{w}_{1,n}\right) \right\} \mu^n \right\} \qquad (52)
\end{aligned}
$$

(51) is the integral formalism of the first kind of independent solution of Lame function for the infinite series. And, (52) is the integral formalism of the second kind of independent solution of Lame function for the infinite series.

5 Application to Laplace equation in ellipsoidal coordinates and nonlinear evolution equations

A great many authors have worked on applications to boundary value problems for the Laplace equation in elliptic coordinates. We can apply the power series expansion in closed forms of Lame function and its integral forms into many mathematical physics areas. For example, Lame equation can be employed to solve boundary-value problems for Laplace equation in elliptical cones. In “Occurrence of periodic Lame functions at bifurcations in chaotic Hamiltonian systems”[3], the authors shows that Lame equation in the Weierstrass’s form occurs at bifurcations in chaotic two-dimensional Hamiltonian systems with mixed phase space. (see (6), (7) in Ref.[3]); Lame equation in the algebraic form can be transforms to in the Weierstrass’s form by changing an independent variable. Then we can describe Lame function more analytically in power series expansions in closed forms and its integral forms. In “Lame Function and Its Application to Some Nonlinear Evolution Equations”[5], the Lame equation is applied to solve nonlinear $(1+1)$-dimensional, $(1+2)$- dimensional and coupled evolution equations. (see (1), (3), (5), (6) in Ref.[5])

6 Conclusion

The power series expansion of Lame equation (ellipsoidal harmonics equation) has a relation between three different coefficients as we see (3). Because of its three term recurrence, there is a mathematical difficulty that Lame functions are described in the power series expansion in closed forms and its integral forms.

In this paper I derive the power series expansion in closed forms for the infinite series and the polynomial of Lame function analytically by applying three term recurrence formula[1]. Even if coefficient α can have two different values in which are $2(2\alpha_i+i+\lambda)$ or $-2(2\alpha_i+i+\lambda)-1$ to make B_n term terminated at certain eigenvalues, its solutions are completely equivalent to each other as we see (7).

Also as we see representations in the form of integrals in Lame functions, a ${}_2F_1$ function recurs in each of sub-integral forms: the first sub-integral form contains zero term of $A_n's$, the second one contains one term of A_n's, the third one contains two terms of A_n's, etc. Then we can transform Lame functions to all other well-know special functions such as Hypergeometric, Bessel, Legendre, Kummer functions, etc.

Since we get the integral forms of power series expansions in Lame function, we are able to obtain generating functions of it. The generating functions are really helpful in order to derive orthogonal relations, recursion relations and expectation values of physical quantities.

Most of well-known papers with boundary value problems in ellipsoidal geometry have been published with using the Weierstrass's form of it. So we need know what the power series expansion in closed forms of Lame function in the Weierstrass's form and its integral forms. In [16], I am deriving the power series expansion in closed forms of Lame function in the Weierstrass's form and its integral forms analytically. In [17], I am investigating the generating functions of Lame equation in the Weierstrass's form.

7 Series “Special functions and three term recurrence formula (3TRF)”

This paper is 6th out of 10.

1. “Approximative solution of the spin free Hamiltonian involving only scalar potential for the $q-\overline{q}$ system” [10] - In order to solve the spin-free Hamiltonian with light quark masses we are led to develop a totally new kind of special function theory in mathematics that generalize all existing theories of confluent hypergeometric types. We call it the Grand Confluent Hypergeometric Function. Our new solution produces previously unknown extra hidden quantum numbers relevant for description of supersymmetry and for generating new mass formulas.

2. “Generalization of the three-term recurrence formula and its applications” [11] - Generalize three term recurrence formula in linear differential equation. Obtain the exact solution of the three term recurrence for polynomials and infinite series.

3. “The analytic solution for the power series expansion of Heun function” [12] - Apply three term recurrence formula to the power series expansion in closed forms of Heun function (infinite series and polynomials) including all higher terms of A_ns.

4. “Asymptotic behavior of Heun function and its integral formalism”, [13] - Apply three term recurrence formula, derive the integral formalism, and analyze the asymptotic behavior of Heun function (including all higher terms of A_ns).

5. “The power series expansion of Mathieu function and its integral formalism”, [14] - Apply three term recurrence formula, analyze the power series expansion of Mathieu function and its integral forms.

6. “Lame equation in the algebraic form” [15] - Applying three term recurrence formula, analyze the power series expansion of Lame function in the algebraic form and its integral forms.

7. “Power series and integral forms of Lame equation in the Weierstrass’s form and its asymptotic behaviors” [16] - Applying three term recurrence formula, derive the power series expansion of Lame function in the Weierstrass’s form and its integral forms.

8. “The generating functions of Lame equation in the Weierstrass’s form” [17] - Derive the generating functions of Lame function in the Weierstrass’s form (including all higher terms of A_n’s). Apply integral forms of Lame functions in the Weierstrass’s form.

9. “Analytic solution for grand confluent hypergeometric function” [18] - Apply three term recurrence formula, and formulate the exact analytic solution of grand confluent hypergeometric function (including all higher terms of A_n’s). Replacing μ and $\varepsilon\omega$ by 1 and $-q$, transforms the grand confluent hypergeometric function into Biconfluent Heun function.

10. “The integral formalism and the generating function of grand confluent hypergeometric function” [19] - Apply three term recurrence formula, and construct an integral formalism and a generating function of grand confluent hypergeometric function (including all higher terms of A_n’s).

References

1. Choun, Y.S., Generalization of the three-term recurrence formula and its applications. arXiv:1303.0806
2. Lame, G., Sur les surfaces isothermes dans les corps homogenes en equilibre de temperature. J. Math. Pures. Appl., 2 (1837), 147-188 (French).
3. Brack, M., Mehta, M. and Tanaka, K., Occurrence of periodic Lame functions at bifurcations in chaotic Hamiltonian systems. J. Phys. A, 34(40) (2001), 8199-8220.
4. Bronski, J. C. , Carr, L. D., Deconinck, B., Kutz, J. N. and Promislow, K., Stability of repulsive Bose-Einstein condensates in a periodic potential. Phys. Rev. A, 3(63) (2001), 1-11.
5. Zun-Tao,F., Shi-Kuo, L., Shi-Da,L. and Qiang1, Z., Lame Function and Its Application to Some Nonlinear Evolution Equations. Commun. Theor. Phys. (Beijing, China), 40 (2003), 53-56.
6. Bateman, H., Erdelyi, A., Higher transcendental functions 3. Automorphic functions. McGraw-Hill, 1955.
7. Hobson, E.W., The theory of spherical and ellipsoidal harmonics. Cambridge Univ. Press, 1931.
8. Whittaker, E.T. and Watson, G.N., A course of modern analysis. Cambridge Univ. Press, 1952.
9. Romain, G. and Jean-Pierre, B., Ellipsoidal Harmonic expansions of the gravitational potential: Theory and application. Celest. Mech. Dyn. Astron., 79(4) (2001), 235–275.
10. Link to arXiv:1302.7309.
11. Link to arXiv:1303.0806.
12. Link to arXiv:1303.0830.

13. Link to arXiv:1303.0876.
14. Link to arXiv:1303.0820.
15. Link to arXiv:1303.0873.
16. Link to arXiv:1303.0878.
17. Link to arXiv:1303.0879.
18. Link to arXiv:1303.0813.
19. Link to arXiv:1303.0819.